

VIEWPOINT

Care Fragmentation, Care Continuity, and Care Coordination—How They Differ and Why It Matters

Lisa M. Kern, MD, MPH
Weill Cornell Medicine,
New York, New York.

Julie P. W. Bynum, MD, MPH
University of Michigan,
Ann Arbor.

Harold Alan Pincus, MD
Columbia University,
New York, New York.

Health care in the US is characterized by fragmentation, with many patients seeing multiple physicians. Indeed, 35% of Medicare beneficiaries saw 5 or more physicians in 2019.¹ Having multiple physicians may be appropriate, but it may also lead to medical errors, unnecessary visits, avoidable hospitalizations, and suboptimal care if all of the physicians do not have complete information about the patient and each other's care plans. Even after widespread dissemination of electronic health records, 34% of primary care physicians in a national study reported that they do not always or most of time receive useful information from specialists about the patients they referred.²

Despite how common it is for patients to see multiple physicians, care fragmentation has received surprisingly little scientific attention. In this Viewpoint, we propose conceptualizing fragmentation as a pattern of health care utilization that could cause harm and that is related to but distinct from care continuity and care coordination. We consider why fragmentation occurs and suggest a potential path forward for developing evidenced-based strategies that can reduce the occurrence of fragmentation and its associated harms.

Definitions

Care fragmentation is care that is diffusely spread across many physicians, such that no single physician accounts for a substantial proportion of visits. Care continuity is the use of the same ambulatory physician repeatedly over time, such that the usual physician (who can be a primary care physician or specialist) accounts for a substantial proportion of visits. Both fragmentation and continuity are measures of utilization. They can both be measured with numerical indices, such as the Bice-Boxerman index (BBI), in which the raw BBI is a measure of continuity and the inverse ($1 - \text{BBI}$) is a measure of fragmentation.^{3,4}

By contrast, care coordination cannot simply be assessed by a quantitative measure of ambulatory utilization. It involves a more qualitative assessment of the extent to which physicians are collectively operating in a teamlike manner to develop and implement an overall care plan to meet the patient's goals.⁵ Effective care coordination can occur in the context of continuous care, fragmented care, both, or neither.

Existing Literature

The continuity end of the ambulatory care utilization spectrum has been studied for decades.⁶ Conceptually, interventions to increase continuity would seek to increase the frequency of returning to the usual provider of care. However, focusing on continuity with a single physician may overlook the other physicians

involved in a patient's care. Care coordination has also been studied for decades, but many interventions designed to improve care coordination have been ineffective.⁷ Effectiveness of coordination interventions may have been limited in part because they have typically focused on a small subset of the complete care team. Meanwhile, there have been relatively few studies of fragmentation and few interventions explicitly designed to address it.

The Need for a Focus on Fragmentation

Why fragmentation occurs in today's health care environment is almost certainly multifactorial, driven by factors ranging from the patient level to the policy level. Thus, the potential solutions will also likely need to be multifactorial. In the **Table**, we summarize potential drivers of fragmentation and potential corresponding interventions, informed by a previous qualitative study⁸ and by the 6P model for developing multilevel strategies.⁹ The 6 levels of the 6P model are the (1) patients (and caregivers), (2) providers (physicians), (3) practice settings, (4) plans (public and private payers), (5) purchasers (employers), and (6) populations (communities).

Highlighting these different levels at which fragmentation develops illuminates different levels at which interventions could be deployed, each with a distinct role in either decreasing fragmentation or mitigating its downstream negative consequences. For example, a patient-level intervention could involve educating and encouraging patients and caregivers to speak up if they notice gaps in communication among physicians. A physician-level intervention could involve (with appropriate plan-based incentives) improving communication among physicians for their common patients and creating shared accountability for patient outcomes. A practice-level intervention could involve creating alerts to notify physicians when they are seeing patients who have many ambulatory physicians and encouraging communication among members of the care team. The list goes on, and the strategies shown in the **Table** are illustrative, not comprehensive. The main point is that looking through the lens of fragmentation illuminates opportunities for intervention that would not have been systematically considered if one were looking through the usual lens of continuity or coordination.

Conclusions

Even though continuity and fragmentation are the mathematical inverses of each other, improving continuity (ie, increasing the number of visits with one physician) would not necessarily decrease the total number of physicians involved in a person's care, nor would it necessar-

Corresponding Author: Lisa M. Kern, MD, MPH, Weill Cornell Medicine, 420 E 70th St, Box 331, New York, NY 10021 (lmk2003@med.cornell.edu).

Table. Generating Ideas for Potential Interventions by Thinking Through Multilevel Drivers of Health Care Fragmentation

| Level of action (6P model) | Contributors to health care fragmentation and subsequent harms | Potential interventions |
|-----------------------------------|--|---|
| Patients (and caregivers) | Assumption that physicians are communicating with each other | Educate and encourage patients and caregivers to bring medical records, test reports, and medicine bottles to visits and report perceived gaps in communication |
| Providers (physicians) | Lack of communication with each other regarding their common patients | With appropriate incentives (see also Plans), improve communication among physicians for their common patients and create shared accountability |
| Practice settings | Lack of notifications for physicians when they are seeing a patient with many ambulatory physicians | Create alerts, notifying physicians when they are seeing patients who have many ambulatory physicians and encouraging communication among members of the care team |
| Plans (public and private payers) | Undervaluing of cognitive services, putting pressure on primary care physicians to see many patients per day | Increase compensation for cognitive services, enabling more time to be spent per patient |
| | Failure to incentivize communication among physicians | Add financial (or other) incentives for primary care physicians and specialists to recognize their shared accountability for patient outcomes, fostering communication and teamlike processes |
| | Lack of use of claims to identify patients with highly fragmented care | Use claims data to identify patients with highly fragmented care and partner with physicians to reduce unnecessary fragmentation |
| Purchasers (employers) | Lack of identification of fragmented care as an opportunity for improvement | Put pressure on public and private payers to hold them accountable for decreasing unnecessary fragmentation |
| Populations (communities) | Lack of consistent communication among community-based urgent care centers and patients' other physicians | Improve communication between urgent care centers and patients' other physicians |

ily improve coordination among all of those physicians. Because the US has not focused on fragmentation, there is no comprehensive inventory of possible interventions that could be deployed to address fragmentation, and it has not yet been determined which interventions are most feasible or most effective. Similarly, there is no

consensus on whose responsibility it is to design, fund, implement, or participate in interventions to address fragmented care. National dialogue and more federally funded research on this issue are urgently needed. Patients are experiencing avoidable harm from fragmented care, and they deserve better.

ARTICLE INFORMATION

Published Online: January 29, 2024.
doi:10.1001/jamainternmed.2023.7628

Conflict of Interest Disclosures: Dr Kern reported personal fees from Mathematica outside the submitted work. Dr Pincus reported personal fees from AbleTo, Magellan Health Studio, and the National Committee on Quality Assurance outside the submitted work. No other disclosures were reported.

Funding/Support: Drs Kern and Bynum were supported by the National Institute on Aging of the National Institutes of Health (award number U54AG063546), which funds the Imbedded Pragmatic Alzheimer's Disease and AD-Related Dementias Clinical Trials Collaboratory. Dr Kern was also funded by the Agency for Healthcare Research and Quality (K18HS029255). Dr Pincus was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health through grant number UL1TR001873.

Role of the Funder/Sponsor: The National Institute on Aging Imbedded Pragmatic Alzheimer's Disease and AD-Related Dementias Clinical Trials Collaboratory reviewed this manuscript to ensure

that their funding was listed accurately but otherwise played no role in the development of this Viewpoint. The other funders played no role in the development of this Viewpoint.

Disclaimer: This content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Agency for Healthcare Research and Quality.

REFERENCES

1. Barnett ML, Bitton A, Souza J, Landon BE. Trends in outpatient care for Medicare beneficiaries and implications for primary care, 2000 to 2019. *Ann Intern Med*. 2021;174(12):1658-1665. doi:10.7326/M21-1523
2. Timmins L, Kern LM, O'Malley AS, Urato C, Ghosh A, Rich E. Communication gaps persist between primary care and specialist physicians. *Ann Fam Med*. 2022;20(4):343-347. doi:10.1370/afm.2781
3. Bice TW, Boxerman SB. A quantitative measure of continuity of care. *Med Care*. 1977;15(4):347-349. doi:10.1097/00005650-197704000-00010
4. Liu CW, Einstadter D, Cebul RD. Care fragmentation and emergency department use

among complex patients with diabetes. *Am J Manag Care*. 2010;16(6):413-420.

5. Care coordination. Agency for Healthcare Research and Quality. Accessed November 8, 2023. <https://www.ahrq.gov/ncepcr/care/coordination.html>

6. Saultz JW, Lochner J. Interpersonal continuity of care and care outcomes: a critical review. *Ann Fam Med*. 2005;3(2):159-166. doi:10.1370/afm.285

7. Peikes D, Chen A, Schore J, Brown R. Effects of care coordination on hospitalization, quality of care, and health care expenditures among Medicare beneficiaries: 15 randomized trials. *JAMA*. 2009;301(6):603-618. doi:10.1001/jama.2009.126

8. Kern LM, Safford MM, Slavin MJ, et al. Patients' and providers' views on causes and consequences of healthcare fragmentation in the ambulatory setting: a qualitative study. *J Gen Intern Med*. 2019;34(6):899-907. doi:10.1007/s11606-019-04859-1

9. Pincus HA, Hough L, Houtsinger JK, Rollman BL, Frank RG. Emerging models of depression care: multi-level ('6 P') strategies. *Int J Methods Psychiatr Res*. 2003;12(1):54-63. doi:10.1002/mpr.142